

Brad S. Elkins, M.D., F.A.C.S.
Postoperative Cataract Care Instructions
(818) 990-3623

1. The day of surgery

- 2-3 hours after you arrive home use your **OFLOXACIN KETOROLAC** and **PREDNISOLONE ACETATE** eye drops.
- Use all 3 drops again at dinner & bedtime.
- Do not press or rub the eye.
- Sleep with the shield tonight.



2. Normal postoperative complaints

- Your eye may be red.
- Your vision may be blurred.
- Your pupil may be large and dilated.
- Your eye may feel scratchy, teary, and light sensitive.
- You may see floaters or black spots.

3. The day after the surgery

- You do not need to wear the shield any more.
- You can resume normal activities.
- Bending and lifting are OK as long as there is no discomfort.
- Showering or bathing are OK but avoid getting water in your eye.
- Do not wear makeup for 1 week.



4. Postoperative eye drops

- OFLOXACIN** 1 DROP 3 X A DAY for 7 days.
- KETOROLAC** 1 DROP 3 X A DAY for 1 month.
- PREDNISOLONE ACETATE** (Shake well) 1 DROP 3 X A DAY for 1 month.
- Wait a few minutes between each drop. **You may finish one bottle before another, this is OK. Just use each as instructed.**
- For instructional video see <http://www.youtube.com/watch?v=-LZw946ahDc>



- 5. Please call the office immediately if there is a serious problem.** Increased redness, pain, swelling, or worsening blurred vision requires an immediate phone call or attention.